



SENSITITRE COMPANION/EQUINE PLATE FORMAT

Plate Code: **COMEQ2F**

Plate Type: **MIC**

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
|----------|-------------|--------------|----------------|-------------|-------------|-----------|-----------|-------------|-------------|-------------|-----------|------------|
| A | SXT 2/38 | SXT 1/19 | SXT 0.5/9.5 | AMP 16 | PEN 8 | CEP 16 | GEN 8 | ENRO 4 | TET 8 | CHL 16 | FOX 16 | ERY 4 |
| B | AMI 32 | AUG 32/16 | TIM 64/2 | AMP 8 | PEN 4 | CEP 8 | GEN 4 | ENRO 2 | TET 4 | CHL 8 | FOX 8 | ERY 2 |
| C | AMI 16 | AUG 16/8 | TIM 32/2 | AMP 4 | PEN 2 | CEP 4 | GEN 2 | ENRO 1 | TET 2 | CHL 4 | FOX 4 | ERY 1 |
| D | AMI 8 | AUG 8/4 | TIM 16/2 | AMP 2 | PEN 1 | CEP 2 | GEN 1 | ENRO 0.5 | TET 1 | TIO 4 | FOX 2 | ERY 0.5 |
| E | AMI 4 | AUG 4/2 | TIM 8/2 | AMP 1 | PEN 0.5 | FAZ 16 | POD 16 | TIC 64 | MAR 2 | TIO 2 | IMI 8 | NEG |
| F | RIF 2 | RIF 1 | AMP 0.25 | AMP 0.5 | PEN 0.25 | FAZ 8 | POD 8 | TIC 32 | MAR 1 | TIO 1 | IMI 4 | POS |
| G | OXA+ 4 | CLI 2 | CLI 0.5 | CLI 0.25 | PEN 0.12 | FAZ 4 | POD 4 | TIC 16 | MAR 0.5 | TIO 0.5 | IMI 2 | POS |
| H | OXA+ 2 | ORB 4 | ORB 2 | ORB 1 | PEN 0.06 | FAZ 2 | POD 2 | TIC 8 | MAR 0.25 | TIO 0.25 | IMI 1 | POS |

ANTIMICROBICS

| | |
|------|--|
| AMI | Amikacin |
| AMP | Ampicillin |
| AUG | Amoxicillin / clavulanic acid 2:1 ratio |
| CEP | Cephalothin |
| POD | Cefpodoxime |
| CLI | Clindamycin |
| SXT | Trimethoprim / sulfamethoxazole |
| ENRO | Enrofloxacin |
| ERY | Erythromycin |
| FAZ | Cefazolin |
| FOX | Cefoxitin |
| GEN | Gentamicin |
| IMI | Imipenem |
| ORB | Orbifloxacin |
| OXA+ | Oxacillin+2%NaCL |
| PEN | Penicillin |
| RIF | Rifampin |
| CHL | Chloramphenicol |
| MAR | Marbofloxacin |
| TET | Tetracycline |
| TIC | Ticarcillin |
| TIM | Ticarcillin / clavulanic acid constant 2 |
| TIO | Ceftiofur |
| POS | Positive Control |
| NEG | Negative Control |